

STATE DEPENDENT δ -PAIRING AND SPONTANEOUS FISSION

A. BARAN, Z. LOJEWSKI and K. SIEJA

*Institute of Physics, University of M. Curie-Skłodowska,
ul. Radziszewskiego 10, 20-031 Lublin, Poland*

Received September 22, 2003

Revised October 30, 2003

We examine the fission barriers, mass parameters and spontaneous fission half lives of fermium isotopes within the framework of macroscopic-microscopic model with a δ -pairing interaction. Four different macroscopic models are applied and studied. The results are compared to experimental data and to the corresponding monopole pairing ones. The half lives obtained in the δ -pairing model are comparable with experimental data. The state-dependence of pairing has an important effect on the calculated fission half-lives.

The calculations of fission half lives T_{sf} in macroscopic-microscopic methods consist of determining the fission barrier (collective potential energy) and the mass tensor of the nucleus undergoing the fission process. The potential energy V splits into a shell δE_{shell} and a pairing correction δE_{pair} (microscopic part) and the smooth average (macroscopic) energy such as liquid drop¹, droplet², LSD³ or Yukawa plus Exponential.⁴

The microscopic part of the energy is calculated using the single-particle Woods-Saxon potential with the *universal* set of parameters.⁵

All the former calculations of T_{sf} were done assuming that the pairing matrix elements $G_{ab} \equiv \langle \nu\bar{\nu} | \hat{V}_{\text{pair}} | \mu\bar{\mu} \rangle$ are constant. Such an approximation leads to the averaging of the superconductive properties of nuclei and weakly reflects the structure of nucleon pairs. In the following we have used the state dependent pairing model where the pairing matrix elements G_{ab} are state dependent and the interaction $\hat{V}_{\text{pair}}$ is of the form:⁶

$$\hat{V}_{\text{pair}} = -V_0 \frac{1 - \sigma_1 \cdot \sigma_2}{4} \delta(\vec{r}_{12}). \quad (1)$$

Here V_0 is a constant determined from experimental nuclear masses.^{7,8}

This form of the interaction better reproduces the properties of the nucleons coupled in pairs in degenerate time reversed states. The δ -pairing strength constants V_0 for protons (p) and neutrons (n) are $V_0^p = V_0^n = 215 \text{ MeV fm}^3$. The masses⁸ in the vicinity of the fermium nuclei are reproduced in this case with an accuracy given by the standard deviation error $\sigma = 0.608 \text{ MeV}$ (see Fig. 1). The residual pairing interaction is treated in the BCS approximation.

Spontaneous fission of a nucleus is described as a tunnelling of the collective potential barrier described in WKB approximation. In order to minimize the ac-

2 *A. Baran, Z. Lojewski and K. Sieja*




Fig. 1. Mass deviations for 47 isotopes of $Z=99, 100, 101$ in the case of the state-dependent δ -pairing and the liquid drop model.

tion which enters the fission probability we have used the dynamic programming method.^{9,10}

The macroscopic-microscopic method is not analytical. Therefore, it is necessary to calculate the potential energy and all the components of the tensor of inertia $B_{kl}(\text{def})$ on a mesh in the multi-dimensional space spanned by a set of collective degrees of freedom consisting of three nuclear shape parameters ($\beta_2, \beta_4, \beta_6$). The collective mass $B_{kl}(\text{def})$ is calculated in the adiabatic cranking model.




Fig. 2. The fission barriers for ^{252}Fm in case of both $G = \text{const}$ (dashed curve) and δ -pairing (solid curve) models.

An example of fission barriers of ^{252}Fm evaluated in both monopole and δ -pairing models is shown in Fig. 2. The monopole pairing barrier ($G = \text{const}$) is denoted by the dashed line and the δ -pairing one is represented by the solid curve. One observes that the monopole pairing model gives a fission barrier similar to that of the δ pairing force model. However, the barrier height is smaller for $G = \text{const}$ pairing.




Fig. 3. The effective mass parameter for ^{252}Fm for two different pairing models, δ (solid) and $G = \text{const}$ (dashed curve) models of pairing.

Figure 3 shows the effective mass parameters taken along the path to fission. One can see the differences of both monopole pairing (dashed curve) and δ -pairing (solid curve) effective inertia.

As was shown the fission barriers strongly depend on the macroscopic energy model.¹¹ To examine this fact in the case of the δ -pairing model we have studied four most frequently used macroscopic models. The choice of the model strongly influences the spontaneous fission half-lives T_{sf} . This is illustrated in Fig. 4 where the results of calculations of the spontaneous fission half-lives for the fermium isotopes are presented. The Figure shows the results for the Yukawa plus exponential macroscopic energy, Lublin-Strasbourg Drop (LSD), the Mayers-Swiatecki drop and for the droplet model. Triangles correspond to the δ -pairing model and the open circles to the monopole pairing one. The experimental data are denoted by plain diamonds. It is seen that the spontaneous fission half-lives considerably depend on the pairing model. The δ -pairing interaction seems to work better with most macroscopic models as compared to the monopole pairing model.

In our comparative study we have neglected the lowest odd-multipolarities usually represented by β_3 and β_5 deformations which, as was shown elsewhere,¹² considerably decrease the fission barriers especially at large quadrupole deformations.

From our investigations one can draw the following conclusions:

- (i) The fission barriers of the δ -pairing force model are similar to that of the classical pairing, however slightly higher.
- (ii) The state dependent δ -type force influences significantly the spontaneous fission half lives. At the same time the isotopic systematics of T_{sf} does not change.
- (iii) The T_{sf} values calculated in different macroscopic models are considerably different and depend on the pairing model used.
- (iv) The investigations of the pairing interaction and the macroscopic energy models should be continued, in order to obtain the most appropriate fission barriers and spontaneous fission half lives.




Fig. 4. Spontaneous fission half-lives (T_{sf}) of fermium isotopes in four models of the macroscopic energy and two models of pairing interactions. The results of the δ pairing model are denoted by plain triangles and the results of the monopole pairing are represented by open circles. Diamonds correspond to the experimental data.

References

1. W. D. Myers, W. J. Świątecki, Nucl. Phys. **81**, 1 (1966), Ark. Fys., **36**, 343 (1967)
2. W. D. Myers, W. J. Świątecki, Ann. Phys. **84**, 186 (1974)
3. K. Pomorski, J. Dudek, Phys. Rev.C **67**, 044316 (2003)
4. H. J. Krappe, J. R. Nix and A. J. Sierk, Phys.Rev. **C20**, 992 (1979).
5. S. Ćwiok, J. Dudek, W. Nazarewicz, J. Skalski and T. Werner, CPC **46**, 379 (1987).
6. S.J. Krieger, P. Bonche, H. Flocard, P. Quentin, M.S. Weiss, Nucl. Phys. **A517**, 275, (1990).
7. A. Baran, K. Sieja, IJTP, 2004, this issue; K. Sieja, A. Baran, Phys. Rev. C **68**, (2003).
8. G. Audi and A. H. Wapstra, Nucl. Phys. **A595**, 409 (1995).
9. A. Baran, K. Pomorski, A. Łukasiak and A. Sobczewski, Nucl. Phys. **A361**, 83 (1981).
10. A. Staszczak, Z. Lojewski, A. Baran, B. Nerlo-Pomorska, and K. Pomorski, Proceedings of the 3rd International Conference, Casta-Papernicka, 30 August, 22 (1996);
11. Z. Lojewski, A. Baran and K. Pomorski, Acta Phys Pol. B **34**, 1801, (2003)
12. Z. Lojewski, A. Staszczak, Nucl. Phys. **A657**, 134, (1999)